



ENACTS

ENHANCED NATIONAL CLIMATE SERVICES

El Niño-Southern Oscillation (ENSO) Rainfall Probability Training

**Training Module
Tanzania
March 09, 2016
Version 1.0**

March 09, 2016



International Research Institute for Climate and Society (IRI), (2016). El Niño-Southern Oscillation (ENSO) Rainfall Probability Training - Tanzania, March 09, Version 1.0. Palisades: IRI.

This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>) and may be adapted or reproduced with attribution to the IRI and for any non-commercial purpose.

Acknowledgements

The ENACTS team, wishes to thank all the individuals and institutions that contributed in the many ways to the preparation of this manual. The shared technical knowledge, experiences, and perspectives have produced a training module that will have a significant positive impact on the capability of strengthening the ENACTS tools in various East African countries.

Special thanks are extended to the IRI staff members who collaborated in the preparation of the different manuals: Dr. Pietro Ceccato, Luz Cervantes, John delCorral, Dr. Tufa Dinku, Igor Khomyakov, Aisha Owusu, Yohana Tesfamariam Tekeste and Dr. Madeleine Thomson.

The preparation of this manual would not have been possible without the support provided by our donors WHO - Global Framework for Climate Services and UK DfID WISER ENACTS and we extend sincere appreciation to our donors.

CONTENTS

1	El Niño-Southern Oscillation (ENSO) Rainfall Probability Training - Tanzania	1
1.1	El Niño-Southern Oscillation (ENSO) Impact in Africa	1
1.2	Overview	2
1.3	Definition	2
1.4	Interpretation	3
1.5	Access	3
1.6	Case Study - Ethiopia	4
1.7	Exercise - Tanzania	5
1.8	Quiz	7
1.9	Summary	8
1.10	Reference(s)	8

EL NIÑO-SOUTHERN OSCILLATION (ENSO) RAINFALL PROBABILITY TRAINING - TANZANIA

1.1 El Niño-Southern Oscillation (ENSO) Impact in Africa

The climate impacts of El Niño (and its counterpart, La Niña) are not uniform across the world or within the African continent. And similar to other parts of the world, the impacts of El Niño or La Niña on rainfall in Africa also vary according to location and season (please refer to Figures 1.1 and 1.2). When sea surface temperatures in the Niño 3.4 region of the equatorial Pacific are unusually high, Southern and Western Africa have a tendency to be anomalously dry during November - February. Additionally, Northeastern Africa generally becomes anomalously dry during its primary and longer rainy season of June - September. The converse is true in these regions during La Niña years.

However, Eastern Africa is different. When an El Niño is underway in this region, the climate has a substantially increased probability of being unusually wet during the secondary and shorter rainy season of October - December, whereas the region's primary and longer rainy season, March - May, is largely unaffected.

At the local level, the impact of ENSO is best assessed using high quality historical data such as the rainfall products available from the ENACTS database and Maprooms.

El Niño and Rainfall

El Niño conditions in the tropical Pacific are known to shift rainfall patterns in many different parts of the world. Although they vary somewhat from one El Niño to the next, the strongest shifts remain fairly consistent in the regions and seasons shown on the map below.




Fig. 1.1: El Niño and Rainfall

La Niña and Rainfall

La Niña conditions in the tropical Pacific are known to shift rainfall patterns in many different parts of the world. Although they vary somewhat from one La Niña to the next, the strongest shifts remain fairly consistent in the regions and seasons shown on the map below.




Fig. 1.2: La Niña and Rainfall

1.2 Overview

Why was it developed?

- The ENSO (El Niño and La Niña) Rainfall Probability Mappings were initially created to help stakeholders identify whether or not rainfall in their country, region, or district of interest was historically associated with ENSO. Where strong relationships exist, rainfall is potentially predictable using ENSO informed seasonal climate forecasts.

What can the ENSO Rainfall Probability Maproom be used for?

- Assessing the historical probability of rainfall for a particular season and within a certain ENSO phase for a specific region or district
- Visualizing the spatial extent of ENSO climate impacts

What can the ENSO Rainfall Probability Maproom NOT be used for?

- Providing information on the current ENSO event

1.3 Definition

ENSO rainfall probability is calculated from the long-term average (1981-2010) of monthly rainfall from the ENACTS rainfall database, then and classified according to ENSO state (El Niño, La Niña, Neutral).

The ENSO state for each season is defined according to the Oceanic Niño Index (ONI). It is calculated using Sea Surface Temperature (SST) anomalies, based on the 1981-2010 normal, in the geographical box (170°W, 5°S, 120°W, 5°N). A season is considered El Niño (La Niña) if it is part of at least 5 consecutive overlapping 3-month long seasons where the ONI is above 0.45°C (below -0.45°C).

1.4 Interpretation

Figure 1.3 is an example of the historical probability of seasonal average monthly rainfall product conditioned on El Niño during the July-August-September season, and falling within the upper (wet) one-third (“tercile”) of the 1983-2010 historical distribution in rainfall for Madagascar. Please note that this is not a forecast.

And also please note that when interpreting the graph, each tick represents the beginning of the year (look at Figure 1.3).




Fig. 1.3: Explanation on the ticks: Each tick on the time axis (x-axis), marks the beginning of the respective year

1.5 Access

The ENSO Rainfall Mapage can be accessed via the Climate Forecast Maproom. http://maproom.meteo.go.tz/maproom/Climatology/Climate_Forecast/ENSO_Prob_Precip.html




Fig. 1.4: Tanzania ENSO Rainfall Forecast Maproom

The Maproom allows you to create an analysis by 3-month period (e.g. Jan-Mar), ENSO state (El Niño, La Niña and Neutral), and rainfall outcome/tercile (wet, normal or dry).

1.6 Case Study - Ethiopia

The Ethiopian climate is extremely variable and complex. Annual rainfall characteristics of Ethiopia are classified into three distinct rainy seasons: (1) the longer, primary season (Jun–Sep: JJAS); (2) the shorter, secondary season (Feb–May FMAM); and (3) the dry season (Oct–Jan: ONDJ). The first season corresponds with the Sahelian rainy season (JAS); whereas, the last two seasons correspond with the main East African seasons (MAM and OND). The seasons are locally defined as Kiremt, Belg, and Bega and respectively. Because Ethiopia’s climate is the most complex topography on the African continent, precise delineation of distinct regions and rainy seasons are difficult as the climate varies significantly within a short distance.




Fig. 1.5: Historical probability of seasonal monthly averages of rainfall conditioned on El Niño in Ethiopia a) low and dry in Jul-Sep (b) high and wet in Oct-Dec

The high resolution ENACTS rainfall database has been used to help determine and associate where and when ENSO impacts rainfall in Ethiopia, as seen in Figure 1.4. However, due to the relative short time series of 1983–2010 and the relatively few ENSO events, it should be noted that the relationships observed may not be statistically significant.

1.6.1 Examples

Figure 1.6 displays the historical impact of ENSO on rainfall via a bar graph product generated after setting the analysis criteria to the following: (1) Region: Ethiopia; (2) Spatially Average Over: Zone, South Wollo, Amhara, Ethiopia; (3) Season: Jul-Sept; (4) Tercile: Dry; and (5) ENSO State: El Niño.

As can be seen, this zone has a strong historical tendency for below normal to normal rainfall during El Niño events (lower tercile/below normal is below black line) and a propensity for above normal rainfall during La Niña events (upper tercile/above normal is above green line). Also, in South Wollo it can be seen that the strong El Niño of 1997 was associated with a July-September drought due to below normal rainfall.

Figure 1.7 displays the historical impact of ENSO on rainfall via a bar graph product generated after setting the analysis criteria to the following: (1) Region: Ethiopia; (2) Spatially Average Over: Zone, Borena, Oromia, Ethiopia; (3) Season: Oct-Dec; (4) Tercile: Wet; and (5) ENSO State: El Niño.

As can be seen, this zone was heavily impacted by the 1997/8 El Niño with extreme, above-normal rainfall. Furthermore, Borena has a tendency to have above normal rainfall during many El Niño years. However, there are notable El




Fig. 1.6: July – September Rainfall Conditioned on El Niño

Niño years when rainfall was normal or below normal.

1.6.2 Case Study Summary

In Ethiopia, the relationship between ENSO and rainfall is significant and varies according to season and region. El Niño years are often associated with major droughts during the July-September primary, rainy season across the central and northern regions of the country, often resulting in food insecurity.

These droughts may then be followed by unseasonal and above normal rains in the southern part of the country that are associated with El Niño impacts on the shorter rainy season of Eastern Africa.

Because ENSO has a strong impact on the magnitude and duration on rains in some seasons and regions, there is the possibility to develop robust and skillful seasonal climate forecasts that may predict the extent of the rains several months in advance. These forecasts may be able to help decision-makers in a number of sectors including agriculture, health, water, energy, disasters etc.

1.7 Exercise - Tanzania

Tanzania has a tropical climate but has regional variations due to topography. Seasonal rainfall is driven mainly by the migration of the Intertropical Convergence Zone. It migrates southwards through Tanzania in October to December, reaching the south of the country in January and February, and returning northwards in March, April, and May. This causes the north and east of Tanzania to experience two distinct rainy seasons – (1) the short rains (Vuli”) from October-December and (2) the long rains (Masika) from March-May. However, the southern, western, and central parts of the country typically only experience one rainy season that continues October-April/May.

Please perform the following exercises:

E1. Proceed to the ENSO Rainfall Probability Mapage within Tanzania’s Climate Forecast Maproom, and set the analysis tool bar to the categories in Figure 1.8. (1) Region: Tanzania; (2) Spatially Average Over: TBC (do not select anything for now); (3) Season: March-May; (4) Tercile: Wet; and (5) ENSO State: El Niño.




Fig. 1.7: October – December Rainfall Conditioned on El Niño



Fig. 1.8: Control Bar Settings for ENSO Rainfall Probability

E2. Now, generate the time series, bar graph for the following provinces: (1) Kasena Nankana East, (2) Central Tongu, (3) Ho West, Volta, (4) and (5) Ruvuma.

E3. For each region, indicate the number of years for the categories in the table below (Figure 1.9). Please note that in the below table and within the time series, bar graph, wet=above normal=above rainfall tercile 0.66667 (green) and dry=below normal=below rainfall tercile 0.33333 (black).

Region	Above Normal	Normal	Below Normal
XXX			
El Nino			
Neutral			
La Nina			

Fig. 1.9: ENSO Phase and Tercile Category Table

1.7.1 Example - Kasena Nankana East District (Figure 1.10)




Fig. 1.10: Time Series Bar Graph and Completed Table for the Arusha Region

1.8 Quiz

Please answer the following questions using the ENSO Rainfall Probability Mapage, and figures and tables generated in the previous exercise

- Q1. In north and east of Tanzania, which rainy season has an increased possibility to be wetter than normal during an El Niño? (a) October-December (b) March-May
- Q2. What year(s) (if any) was consistently the wettest for all regions examined for Oct-Dec (OND) and associated with El Niño?
- Q3. Which region(s) had the highest, above-normal rainfalls associated with El Niño for OND?

Q4. How do the magnitudes of rainfall associated with El Niño events in Tanzania OND compare with that of Ethiopia OND? (a) Weaker (b) Same (c) Stronger

1.8.1 Quiz - Answers

A1. October-December

A2. 1986, 1997, 2006

A3. Arusha, Kigoma, Ruvuma

A4. (b) Same

1.9 Summary

ENSO impacts the climate of Tanzania during the primary rainy season of October-December, especially within the northern and eastern regions. During El Niño the rains tend to increase whereas during La Nina they decrease. Temperatures are also affected by El Niño and La Niña.

1.10 Reference(s)

- Kousky, V. E. and Higgins, R. W. (2007). An Alert Classification System for Monitoring and Assessing the ENSO Cycle. *Wea. Forecasting*, 22, 353-371. doi: <http://dx.doi.org/10.1175/WAF987.1>